Research Article

Saraswati Panchak – A Novel Herbal Combination for Mental Health

Ruchi Singh^{1*}

¹Quality Control Head, Shantikunj Pharmacy, Shantikunj, Haridwar, India *Corresponding. Email: <u>ruchi.singh@dsvv.ac.in</u> <u>https://doi.org/10.36018/ijyr.v3i2.59</u>

Abstract. Herbal medicines have a history of long therapeutic use and are still serving a lot of the health needs of a big population of the world. However, the quality control still remains a challenge because of the variation of chemical constituents involved. There are numerous compounds in herbal drugs that are in complex matrices in which no single active constituent is responsible for the overall efficacy. This creates a challenge in establishing quality control standards and standardization of finished herbal drugs. Saraswati Panchak is a novel combination given by Pandit Shri Ram Sharma Acharya for mental health. In this study, components of this novel herbal combination were explained for its potential utility in mental illness. In addition, the study also detailed how Saraswati Panchak Churna was converted into tablet form to increase the shelf life, make it easy to dispense, for dose fixation, etc. The tablet was subjected to organoleptic analysis and physico-chemical analysis to establish its efficacy. After performing complete analysis it was found that all the parameters are within the range. The coarse powder of these herbs may be used as havan samagri in brain related disorders because of the presence of nootropic herbs.

Keywords. Saraswati Panchak, quality control, tablet, Yagya therapy



Introduction

According to World Health Organization mental health is a "state of well-being whereby individuals recognize their abilities, are able to cope with the normal stresses of life, work productively and fruitfully, and make a contribution to their communities." Although mental disorders reflect psychiatric disturbance, people may be affected by mental health problems (1). Mental illness accounts for about one-third of the world's disability caused by all adult health problems, resulting in enormous personal suffering and socioeconomic costs (2). Severe mental health problems including major depressive disorder, bipolar disorder. schizophrenia and substance use disorders affect all age groups and occur in all countries. Mental illness is closely associated with poverty, wars, and other humanitarian disasters, and in some cases, leads to suicide, one of the most common causes of preventable death among adolescents and young adults (3).

A visionary seer sage Pt. Shri Ram Sharma Acharya long back ago realized the need of hour and given a ayurvedic combination named "Saraswati Panchak" to combat with above mentioned problems as the role of herbal medicine in the treatment of various psychological disorders has become well established over the past decade with phytotherapeutic preparations such as St John's wort and kava possessing respectable clinical evidence (4).

Saraswati Panchak comprises of two words, 'Saraswati' and 'Panchak'. The word Saraswati is the name of a goddess, and also symbolizes 'intellect'. Panchak denotes a drug comprises of five elements. Saraswati Panchak is a novel herbal combination containing combination of drugs which are known for potential multiple benefits including mental illness and in psychological disorders. It contains five herbs– Brahmi, Shankhpushpi, Vach, Gorakhmundi and Satavari. The combination gives the properties such as 'brain' tonic, memory enhancer and general health tonic. Brahmi, Vacha, and Shankhapushpi are known Medhya Rasayana or nootropic herbs.

Nootropics are known as memory enhancers, brain tonics and cognitive enhancers. A cognitive enhancer is a substance that enhances memory and concentration. Nootropics are thought to work by altering the availability of the supply of neurochemicals (neurotransmitters, hormones and enzymes) in brain, by stimulating nerve growth or by improving the brain's oxygen supply. Memory herbs are responsible to increase the level of neurotransmitters, particularly acetylcholine and improve blood flow to the brain, by increasing its oxygen and nutrient supply, which will aid memory and brain function (5).

This study presented the overview of the novel combination, its ayurvedic and modern mechanistic inside based on the literature, and its applications in the form of tablet and as a Hawan Samagri (for Yagya Therapy) as well as standardization of preparation of Saraswati Pnachak Vati (tablet) in the herbal pharmacy.

Herbs used in Saraswati Panchak

Brahmi (Bacopamonnieri)

B. monnieri is a creeping perennial with small oblong leaves and purple flowers, found in warm wetlands, and native to Australia and India. It is commonly found as a weed in rice fields, the plant grows throughout East Asia and the United States (6). The whole plant is used medicinally. *B. monnieri* was initially described



around the 6th century A.D. in texts such as the Charaka Samhita, Atharvaved, and Susrutu Samhita as a medhya rasayana–class herb taken to sharpen intellect and attenuate mental deficits. The herb was allegedly used by ancient Vedic scholars to memorize lengthy sacred hymns and scriptures (7).

Brahmi contains alkaloid nicotinine, herpestine, brahmine, bacosides A and B, saponins A, B and C, triterpenoidsaponins, β -sitosterol, stigmastanol, betulinic acid, D-mannitol, stigmasterol, α -alanine, glutamic acid, aspartic acid, and serine (8).

It is used traditionally to treat various nervous disorders, digestive aid, improve learning, concentration and memory to provide relief to patients with anxiety, and skin diseases; specific uses include the treatment of insanity, asthma and epilepsy (9–11).

The Bacopa herb, also called nootropic herb, helps in the repair of damaged neurons, neuronal synthesis, the restoration of synaptic activity, and improves brain function. Numerous studies suggested that R monnieri's bioactive components (ie, bacosides) protect the brain against oxidative damage and age-related cognitive deterioration with several mechanisms of action (12-13), the bioactive constituent, bacoside A was present in the B. monnieri extract (BME)-treated rat serum and could directly or indirectly interact with the neurotransmitter systems to improve memory and learning ability (14). A clinical study also suggested that the oral treatment with B. monnieri was able to enhance memory in both adults and children (15).

Shankhpushpi (Convolvulus pluricaulis)

C. pluricaulis is a perennial herb that seems like morning glory. Its branches are spread on the ground and may be more than 30 cm long. The flowers are blue in color and the leaves, which are elliptical in shape, are located at alternate positions with flowers or branches. Known asAloe weed in English, the herb is commonly found in India, especially in the state of Bihar (16).

C. pluricaulis contains D-glucose, maltose, rhamnose, sucrose, shankhapushpine, convolamine, convoline, convolidine, convolvine, confoline, convosine, volatile oils, fatty acids, fatty alcohols, hydrocarbons, myristic acids, palmitic acids and linoleic acids, Scopoletin, β -sitosterol, ceryl alcohols, 20oxodotriacontanol, tetratriacontanoic acids, flavonoid-kampferol, steroids-phytosterols (17).

Shankhpushpi is a reputed drug of Ayurveda and reported as a brain tonic, nervine tonic, alternative and laxative (18-19). This plant is reported to be a prominent memory improving drug, a psycho-stimulant and tranquilize, and reduce mental tension. There are many references in Ayurvedic texts about the use of this drug as brain tonic in hypotensive syndromes (20).

Two important forms of synaptic plasticity known to be involved in the processes of memory formation are long-term depression (LTD) and long-term potentiation (LTP). In a study, the effect of C. pluricaulis plant extract on LTP and LTD were evaluated and found that prolonged treatment of C. pluricaulis extract, at a specific dose in healthy animals, can augment memory functions by modulating hippocampal plasticity (21).



Vach (Acoruscalamus)

A.calamus (L.) (Sweet flag), a member of the family Acoraceae, generally used alone or in combination with other herbs in Indian and Chinese traditional medicine has generated great interest and is found to be beneficial (22). The plant is widely cultivated in different parts of sub-temperate and temperate regions of the world and is native to India, Japan, Sri Lanka, China, Mongolia, Burma, Europe, SouthernRussia and Northern USA (23).

The oil contains varying concentrations of αasarone. β-asarone. γ -asarone. calamenenolcalamene, calameone, camphene, α -pinene, β -pinene, p-cymene, eugenyl acetate, eugenol. isoeugenol, methyl isoeugenol, azulene. calamol, eugenol methyl ether, dipentene, methyl eugenol, asaronaldehyde, terpinolene, camphor, 1,8-cineole, αcaryophyllene and hydrocarbons (24-26). The oil also contains fatty acids such as palmitic acid and its ester, heptylic acid, an ester of butyric acid (27).

The traditional use *of A.calamus* in Indian Ayurvedic system is widely accepted. The plant has been used to cure many diseases like cough, fever, asthma, hysteria, epilepsy, depression, insomnia, mental retardation, skin diseases, haemorrhoids, dysentery, diarrhea, kidney and liver problems, bronchitis and as a sedative (28).

Methanol extracts of *A. calamus* showed significant acetylcholine esterase enzyme inhibition at a concentration 200 mg/mL (25). Houghton and coworkers reported the in- vitro acetylcholinesterase inhibitory effect of β asarone and α -asarone. The AChE-inhibitory activity of the oil may be ascribed to β -asarone. Because memory and cognitive performance are related to acetylcholine levels, the AChEinhibitory effect of this plant may account for its traditional use (29).

Gorakhmundi (Sphaeranthusindicus)

S. indicus is a multi-branched herb having round purple flowers which grows plentifully in rice fields and is distributed throughout India, Malay, Ceylon, China and Africa. It is distributed throughout India, Sri Lanka, Africa and Australia from sea level to 1200 m altitude (30).

numbers of phytochemicals Large were isolated from whole plant, leaves and flowers of S. indicus. Aerial parts of this plant showed presence of anessential oil, glycosides, and eudesmanoids, analkaloids phaeranthine and an isoflavone 5,4'-dimethoxy-3'-prenylbiochanin 7o-β-galctosidewith some interesting sesquiterpene, n-pentacosan, hentriacontane, ntriacontanol, β-sitosterol, stigmasterol, β-Dglucoside of β -sitosterol, sphaeranthine and a phenolic glycoside. The essential oil, obtained by steam distillation of the whole herb, contains ocimene, α -terpinene, methyl-chavicol, α -citral, geraniol, α -ion one, β -ionone, d-cadinene, pmethoxy-cinnamaldehyde (31). All the parts of the S. indicus have medicinal uses.

In Ayurvedic system of medicine, the entire plant is used in tuberculous glands, insanity, indigestion, bronchitis, elephantiasis, anaemia, pain in the vagina and uterus, piles, biliousness, epileptic convulsions, leukoderma, dysentery, vomiting, urinary discharges, looseness of the breasts, hemicranias (32).

S. indicus flower derived constituents exhibits synergistic effect against acetylcholinesterase and possess potential antiamnestic activity (33).



Satavari (Asparagus racemosus)

A. racemosus grows throughout the subtropical and tropical parts of India up to an altitude of 1500 m. The plant is a spinous under-shrub, short rootstock bearing with. numerous succulent tuberous roots which are ash colored or silvery white externally and white internally. These roots are use in various medicinal preparations. The stem is climbing, woody, whitish grey or brown colored with small spines. The plant flowers duringFebruary-March leaving a mild fragrance in its surrounding and by the end of April, fruits can be seen with attractive red berries (34).

The powdered roots contain protein, saponins, carbohydrate and oil. The major active chemical constituents of *Asparagus* are a group of steroidal saponins. The plant also contains vitamins A, B1, B2, C, E, P, Mg, Ca, Fe, and folic acid. Other primary chemical constituents of *Asparagus* are essential oils, asparagine, arginine, tyrosine, flavonoids (kaempferol, quercetin, and rutin), resin, and tannin (35).

A. racemosus is a well-known Ayurvedic rasayana which prevents ageing, increase

longevity, impart immunity, improve mental function, vigor and addvitality to the body and it is also used in nervous disorders, dyspepsia, tumors, inflammation, neuropathy and hepatopathy (36).

Methanolic extract of *A. racemosus* showed significant antidepressant-like activity almost certainly by inhibiting Monoamine oxidase (MAO-A and MAO-B) activity; and through interaction with adrenergic, dopaminergic, serotonergic and GABAergic systems (37).

Quality control parameters of Saraswati Panchakvati (tablet)

Procurement of plant material

The crude drugs for the preparation of SP were taken from the pharmacy of Shantikunj, Haridwar after proper authentication. Then, thephysical impurities were removed and the drugs were washedwith water and sun dried below 45°C. Dried drugs were stored in tightly closed containers. The crude drugs used in Saraswati Panchak with their botanical identities and partsused are given in Table 1.

Drug Name	Botanical Name	Family	Part Used	
Brahmi	BacopaMonnieri	Plantaginaceae	Whole Plant	
Shankhpushpi	Convolvulus Pluricaulis	Convolvulaceae	Whole Plant	
Vach	Acoruscalamus	Acoraceae	Root	
Gorakhmundi	SphaeranthusIndicus	Asteraceae	Whole Plant	
Shatawar	Asparagus Racemosus	Asparagaceae	Root	

Table 1. Components of 'Saraswati Panchak' combination



Sr. No.	Test Parameters	<u>Batch</u> <u>No. 1</u>	Batch No.2	Batch No.3	Batch No.4	Batch No.5	Batch No.6	Batch No.7	<u>Batch</u> <u>No.8</u>	Limits
1	Description	Description Brown colored, uncoated biconvex tablets with specific odour and bitter in taste								
2	Weight variation (Table 3)	Within Limits	Within Limits	Within Limits	Within Limits	Within Limits	Within Limits	Within Limits	Within Limits	±5%
3	Hardness of tablet	3.14	3.5	3.9	3.8	3.6	3.1	3.5	3.63	3-6 kg
4	Tablet disintegration time	15 min	12.9 min	15.7 min	14.2 min	15.5 min	13.8 min	13.3 min	14 6 min	NMT 15 min.
5	Friability Test	0.8	0.91	0.95	0.89	0.87	0.92	0.9	0.93	NMT 1%
6	pH of 1% aqueous solution	7.02	7.08	7.06	7	7.06	7.03	7.05	7.06	NA
7	Loss on drying at 105°c (%w/w)	6.36	6.32	6.52	6.45	6.08	5.31	3.16	1.83	NA
8	Total Ash (%w/w)	10.89	11.75	10.52	11.38	12.48	18.38	12.74	12.92	NA
9	Acid Insoluble Ash (%w/w)	2.4	2.4	2.39	2.34	2.44	3.61	0.92	2.12	NA
10	Alcohol Soluble Extrative (%w/w)	3.29	3.26	3.26	3.21	3.28	13.11	19.25	18.31	NA
11	Water Soluble Extrative (%w/w)	38.86	31.78	40.58	37.76	39.58	43.47	39.12	43.06	NA
			Test F	or Heavy M	etals: By AA	AS (all units	s in ppm)	1	1	
12	Lead as Pb Mercury as Hg Arsenic as As Cadmium as Cd	6.8 ND* ND* ND*	ND* ND* ND* ND*	ND* ND* ND* ND*	ND* ND* ND* ND*	5.2 ND* ND* ND*	0.76 ND* ND* ND*	ND* ND* ND* ND*	ND* ND* ND* ND*	NMT 10 NMT 01 NMT 03 NMT 0.3
			Test fo	or Aflatoxins	: By LCMSI	MS (all units	s in ppm)			
13	B1 G1 B1 G2	ND* ND* ND* ND*	ND* ND* ND* ND*	ND* ND* ND* ND*	ND* ND* ND* ND*	ND* ND* ND* ND*	ND* ND* ND* ND*	ND* ND* ND* ND*	ND* ND* ND* ND*	NMT 0.5 NMT 0.5 NMT 0.1 NMT 0.1
14			P	arameter te	sted for Pe	sticide Resi	due	I		
14.1	Azinophos-methyl	ND*	ND*	ND*	ND*	ND*	ND*	ND*	ND*	0.04
14.2	Diazinon	ND*	ND*	ND*	ND*	ND*	ND*	ND*	ND*	0.04
14.3	Dichlorvos	ND*	ND*	ND*	ND*	ND*	ND*	ND*	ND*	0.04
14.4	Malathion	ND*	ND*	ND*	ND*	ND*	ND*	ND*	ND*	0.04
14.5	Methidathion	ND*	ND*	ND*	ND*	ND*	ND*	ND*	ND*	0.04
14.6	Parathion	ND*	ND*	ND*	ND*	ND*	ND*	ND*	ND*	0.04
14.7	Parathion Methyl	ND*	ND*	ND*	ND*	ND*	ND*	ND*	ND*	0.04
15	BIOLIGICAL TESTING AYUSH PRODUCT									
	Test For Microbiology (Total count unit in (cfu/gm)									
15.1	Bacteria Yeast &Mould	3200 620	33000 620	500 29	1570 220	680 <10	54000 200	24x10 ³ 2x10 ²	500 100	1X10⁵ 1X10³
Any Specific Pathogens (unit in (cfu/gm)									·	
15.2	E.coli/gm Salmonella/gm S. aureus/gm P. aeruginosa/gm	Ab Ab Ab Ab	Ab Ab Ab Ab	Ab Ab Ab Ab	Ab Ab Ab Ab	Ab Ab Ab Ab	Ab Ab Ab Ab	Ab Ab Ab Ab	Ab Ab Ab Ab	NA

 Table 2. Analysis of Saraswati Panchak Tablet (Vati). NA= Not Available as it is a new combination; ND=Not Detected; NMT=Not

 More Than; Ab=Absent



Batch No	1	2	3	4	5	6	7	8
Sample (readings)	9	8	12	17	8	10	12	12
Mean (20 tablets weight in grams)	13.02	13.05	13.01	13.13	13.01	13.06	13.07	13.18
Standard Deviation	0.02	0.05	0.05	0.21	0.05	0.05	0.07	0.30
% variation	0.14	0.41	0.08	0.99	0.10	0.45	0.51	1.37

Table 3. Weight Variation Test of Saraswati Panchakvati

Preparation of Saraswati Panchak tablet

All the authenticated crude drugs were crushed to a coarse powder separately and after that mixed thoroughly with eight parts of water in a stainless steel container. Then continuous mild heat was applied until it was reduced to onefourth of its initial quantity. During heating, continuous stirring was done to avoid any deterioration due to burning of materials. After a desirable reduction in volume was achieved, the decoction was filtered through single folded cotton cloth and collected in a separate vessel.Then, the decoction was boiled again with addition of 10% accasia gum over slow fire on a gas stove, maintaining the temperature between 90°C and 95°C till a semisolid consistency is obtained. As the water evaporates he viscosity of the extract increases resulting in 'Ghana' form (38). Then, the Ghana was mixed with the Churna of Saraswati Panchak (up to 10% of extract) further forming a solid mass.Now, the dissolved starch is mixed with the complete mixture. The solid mass (Ghana) was forced through a no. 16 sieve and granules were prepared and then dried at 50°C in a hotair oven for 16 hours. The dried granules were passed againthrough a no. 20 sieve. The combination was then compressed in a single-punch tablet press with a target weight of 650 mg.

Analysis of the Tablet as per Ayurvedic pharmacopoeia of India (API)

Saraswati Panchak tablet were subjected to various analytical parameters (Table 2) as

follows: 1) Organoleptic parameters: Rupa (color), Rasa (taste), Gandha (odor) and Sparsha (touch), 2) Physico-chemical parameters: pH of 5% aqueoussoln., loss on drying at 110°C, ash acidinsoluble value. ash, water soluble extractive. methanolsoluble extractive. 3) Quantitative tests for tablet: Weight variation test (Table 3), tablehardness test, tablet disintegration time, friability, 4) Toxicological: Heavy metal analysis, pesticide residue value, 5) Microbial overload: Bacterial and fungal growth studywas carried out, 6) Reagents and chemicals: All the reagents and chemicalsused for the study were of analytical grade.

Application of Saraswati Panchak coarse powder in Yagya Therapy

Yagya Therapy or Yagyopathy is an ancient Indian method of herbal inhalation therapy that allows for the pulmonary administration of herbs. In Kashyap Samhita an entire chapter named 'DhoopaKalpa' is dedicated on this subject. Various types of 'Dhoopan' are widely used in Ayurvedic treatments for various disorders. During this process, medicinal fume are generated by fumigating the desired herbs or minerals and patient is asked to inhale medicinal herbal fumes, which contains phyto-constituents that provide therapeutic advantages (39-40). The method of medicinal fume inhalation is not a new concept; rather, it prevailed in many traditional and cultural practices for pulmonary and neuronal diseases (41). Herbal fumes produced at high temperature are considered as a



simple way of administering a drug, which exhibits rapid pharmacological activity when inhaled. The herbal medicine combination inhaled in this therapy first reach the brain, followed by lungs and other subtle components of the body. Thus, Yagya therapy has direct biochemical healing effect on CNS tissue diseases & complexities. Acharyas also described multiple drug formulae through nasal route particularly for CNS disorders. Ayurveda also recommended nasal route as a preferred mode of administration of drugs for epilepsy (42).

Now, Yagya Therapy has been proved as a tool for medicinal herbal fumes inhalation as it has shown therapeutic advantage in mental health (43) and epilepsy (44), Saraswati Panchak coarse powder can be used to cure brain related problems (45) as the specific energy currents reduced by Dhoopan and mantra can have significant remedial effect on the disorders & diseases ranging from headache, migraine, intellectual deficiency, depression, seizure disorders (46).

Conclusion

It is now widely accepted that the nature has best answer to all the diseases affecting the human body from time to time. This study has included the Ayurvedic treatments, which is described for mental disorders and are currently part of the Ayurvedic prescriptions. Thus, it could be concluded that the Ayurvedic system of herbal medicine is certainly a treasury of plant drugs for treatment of mental disorders which needs to be explored.

References

- 1. World Health Organization.Invest in mental health. Geneva: WHO, 2003.
- Anderson P, Jane-Llopis E, Hosman C. Reducing the silent burden of impaired mental health. Health Promot Int. 2011;26 (1):4-9. DOI: https://doi. org/10.1093/heapro/dar051. https://doi.org/10.1093/heapro/dar051
- Lake J, Turner MS. Urgent Need for Improved Mental Health Care and a More Collaborative Model of Care. Perm J. 2017;21:17-024. doi: 10.7812/TPP/17-024. https://doi.org/10.7812/TPP/17-024
- Singh YN. Potential for interaction of kava and St. John's wort with drugs.J Ethnopharmacol.2005;100(1-2):108-13.doi: 10.1016/j.jep.2005.05.014. https://doi.org/10.1016/j.jep.2005.05.014
- Amin H, Sharma R. Nootropic efficacy of SatvavajayaChikitsa and Ayurvedic drug therapy: A comparative clinical exposition.Int J Yoga.2015;8(2):109-16.doi: 10.4103/0973-6131.158473.

https://doi.org/10.4103/0973-6131.158473

- Barrett SC, StrotherJL.Taxonomy and natural historyofBacopa in California.Syst Bot. 1978;5:408-419. <u>https://doi.org/10.2307/2418753</u>
- AguiarS,Borowski T. Neuropharmacological review of the nootropic herb Bacopamonnieri.Rejuvenation research. 2013;16(4): 313-326. https://doi.org/10.1089/rej.2013.1431
- Devishree RA, Kumar S, Jain AR. Short term effect of Bacopamonnieri on memory- A brief review.J. Pharm. Res. 2017;11:1447-1450.
- Jyoti A, Sharma D. Neuroprotective role of Bacopamonniera extract against aluminium-induced oxidativestress in the hippocampus of rat brain. NeuroToxicol.2006; 27: 457-7. https://doi.org/10.1016/j.neuro.2005.12.007
- 10. Calabrese C, Gregory WL, Leo M, Kraemer D, Bone K, Oken B. Effects of a standardized Bacopamonnieriextract on cognitive performance, anxiety, and depression in the elderly: A randomized, double-blind,placebo-controlled trial. J. Altern. Complement. Med. 2008 ;14: 707-713. https://doi.org/10.1089/acm.2008.0018
- Kamkaew N, Scholfield CN, Ingkaninan, K,Taepavarapruk N, Chootip K. Bacopamonnieriincreasescerebral blood flow in rat independent of blood pressure.Phytother.Res. 2013; 27: 135-138. <u>https://doi.org/10.1002/ptr.4685</u>
- Saraf MK, Prabhakar S, Khanduja KL, Anand A. Bacopamonnieraattenuatesscopolamine-induced impairment of spatial memory in mice. Evid Based ComplementAlternat Med. 2011;2011:236186.



https://doi.org/10.1093/ecam/neg038

- 13. Mukherjee S, Dugad S, Bhandare R. Evaluation of comparative free-radicalquenching potential of Brahmi (Bacopamonnieri) and Mandookparni (Centellaasiatica). 2011;32:258-264. Ayu. https://doi.org/10.4103/0974-8520.92549
- 14. Rajan KE, Preethi J, Singh HK. Molecular and functional characterization of Bacopamonniera: a retrospective review. Evid Based Complement Alternat Med.;2015:945217.

https://doi.org/10.1155/2015/945217

15. Abdul Manap A S, Vijayabalan S, Madhavan P, Chia YY, Arya A, Wong E H, Rizwan F, Bindal U, KoshyS.Bacopamonnieri, a Neuroprotective Lead in Alzheimer Disease: A Review on Its Properties, Mechanisms of Action, and Preclinical and Clinical Studies. Drug target insights, 2019;13: 1177392819866412.

https://doi.org/10.1177/1177392819866412

- 16. Kumar DC.Pharmacognosy can help minimize accidental misuse of herbal medicine. CurrSci.2007; 93: 1356-1358.
- 17. Agarwal P, Sharma B, Fatima A, Jain SK.An update herb Convolvulus on Ayurvedic pluricaulisChoisy.Asian Pacific journal of tropical biomedicine. 2014;4(3): 245-252. https://doi.org/10.1016/S2221-1691(14)60240-9
- 18. Panda S, Kar A. Inhibition of T3 production in levothyroxine-treated female mice by the root extract of Convolvulus pluricaulis. HormMetab Res. 2001;33(1):16-18.

https://doi.org/10.1055/s-2001-12620

- 19. Sairam K, Rao CV, GoelRK.Effect of Convolvulus pluricaulisChoisy on gastric ulceration and secretion in rats.Indian J Exp Biol. 2001;39(4):350-354.
- 20. Sethiya NK, Nahata A, Mishra SH, Dixit VK.An update on Shankhpushpi, a cognition-boosting Ayurvedicmedicine.Zhong Xi Yi Jie He Xue Bao.2009 ;7(11):1001-22. doi: 10.3736/jcim20091101. PMID: 19912732. https://doi.org/10.3736/jcim20091101
- 21. DasR, SenguptaT, Roy S, Chattarji S, Ray J. Convolvulus pluricaulis extract can modulate synaptic plasticity in rat brain hippocampus.Neuroreport. 2020;31(8):597-604. doi: 10.1097/WNR.00000000001446.

https://doi.org/10.1097/WNR.00000000001446

22. Rajput SB.An overview on traditional uses and pharmacological profile of Acoruscalamus Linn. (Sweet flag) and other Acorus species, Phytomed: Int. J. Phytother. Phytopharm. 2014; 21 (3): 268-276.

https://doi.org/10.1016/i.phymed.2013.09.020

- 23. BalakumbahanR.Acoruscalamus: an overview, J. Med. Plants Res. 2010; 4 (25):2740-2745.
- 24. Nigam MC, Ahmad Α, Misra LN.GC-MS examinationof oil the essential of Acoruscalamus.IndianPerfum.1990; 34: 282-285.
- 25. SrivastavaVK. SinghBM, NegiKS, Pant KC. SunjeaP.Gas chromatographic examination of some aromatic plants of Uttar Pradesh hills. Indian Perfum.1997; 41: 129-139.
- 26. Das BK, ViswanathaSwamyAHM,Koti BC, Gadad, PC.Experimental evidence for use of Acoruscalamus (asarone) for cancer chemoprevention.Heliyon.2019;5(5):e01585. https://doi.org/10.1016/j.heliyon.2019.e01585
- 27. Oh MH, Houghton PJ, Whang WK, Cho JH.Screening of Korean herbal medicines used to improvecognitive anti-cholinesterase function for activity.Phytomedicine.2004; 11: 544-548. https://doi.org/10.1016/j.phymed.2004.03.001
- 28. Chaudhury SS, Gautam SK, Handa KL. Composition of calamus oil from calamus roots growing in Jammu and Kashmir. Indian J Pharm. 1957;19: 183-186.
- 29. Houghton PJ, Kumar V, Govindarajan R, Mukherjee PK.Asarones in Acoruscalamus and theiracetylcholinesterase inhibition. J Pharm PharmacolSupplement.2006;58:A55-A55.
- 30. Rana JS, ReddyMN.Screening for antibacterial activity of SphaeranthusIndicus L. against uropathogenicbacteria.International Journal of Health and Pharmaceutical Sciences.2020,1(1),96-101.
- 31. Bhutkar PM, Suganthi V, Bhutkar MV. Medicinal Uses of SphaeranthusIndicus: A Review. National Journal of Basic Medical Sciences. 2018;8(3):160-166. https://doi.org/10.31975/NJBMS.2018.8307
- 32. Ambavade SD, MhetreNA, TateandVD,BodhankarS L. Pharmacological evaluation of the extracts of Sphaeranthusindicus L. Flowers on anxiolytic activity in mice. Indian J. Pharmcol.2006; 38:254-259.

https://doi.org/10.4103/0253-7613.27021

- 33. Patel MB, Amin D. Sphaeranthusindicus flower derived constituents exhibits synergistic effect against acetylcholinesterase and possess potential antiamnestic activity. J Complement Integr Med. 10.1515/1553-3840.1618. 2012;9(1):doi: PMID: 23023564. https://doi.org/10.1515/1553-3840.1618
- 34. Anonymous. The Wealth of India, Raw materials. Publication and InformationDirectorate, CSIR, N. Delhi.1987;468-472.
- 35. Negi J S. Singh P.JoshiGP.Rawat MS. BishtVK.Chemical constituents of Asparagus.Pharmacogn Rev. 2010;4(8): 215-220. doi: 10.4103/0973-7847.70921 PMCID: PMC3249924 PMID: 22228964.



PUBLISHED BY DEV SANSKRITI VISHWAVIDYALAYA, HARIDWAR, INDIA

https://doi.org/10.4103/0973-7847.70921

36. Alok S, Jain S K, Verma A, Kumar M, Mahor A, Sabharwal M. (). Plant profile, phytochemistry and pharmacology of Asparagus racemosus (Shatavari): A review. Asian Pacific Journal of Tropical Disease.2013;3(3): 242-251.

https://doi.org/10.1016/S2222-1808(13)60049-3

- 37. Dhingra D, Kumar V. Pharmacological evaluation for antidepressant-like activity of Asparagus racemosus wild in mice.Pharmacology online.2007; 3:133-52.
- AcharyaYadavavjiTrikamji.Siddha Yoga Sangraha.
 11th ed. Nagpur: Baidyanath Ayurveda Bhavana;
 2000. p. 4
- Joshi RR, Raghuvanshi M, Pandya P. Yagyopathy versus oral and iv drug administration: evaluation for pulmonary tuberculosis using compartment modeling. J Biol Syst. 2006;14(03):463-89. Available from: http://www.worldscientific.com/doi/abs/10.1142/S021 8339006001891

https://doi.org/10.1142/S0218339006001891

40. Patel V, Mishra A, Shrivastav V. Pulmonary inhalation of medicinal smokes- an aspect of Yagya therapy: an effective therapeutic application and efficient drug delivery model of multiple herbs. In: National Medicinal Plants Board (Ministry of AYUSH, GovtOf India) sponsored National conference on Recent Advances in Ayurvedic Herbal Medicine -Dehradun. 2017:16th Sept Mohagheghzadeh A, Faridi P, Shams-Ardakani M, Ghasemi Y. Medicinal smokes. J Ethnopharmacol. 2006;108(2):161-84

https://doi.org/10.1016/j.jep.2006.09.005

- 42. Jain S, TandonPN.Ayurvedic medicine and Indian literature on epilepsy.Neurol Asia. 2004;9(1):57-8. Available from: <u>http://www.neurology-asia.org/articles/20043_057.pdf</u>
- Sharma S. YagyaChikitsaDwaraManasikSwasthya Par Padane Vale PrabhavKaAdhyayan (Hindi) (Dissertation).DevSanskritiVishwavidyalaya, Shantikunj, Haridwar, Uttarakhand, India; 2009
- 44. Mishra A, BathamL, VermaS, MishraS, ShrivastavaV.Management of Epileptic Seizures through an Integrated Approach including YagyaTherapy, Interdisciplinary Journal of Yagya Research. 2019; 1(2): 52-64.

https://doi.org/10.36018/ijyr.v2i1.24

- 45. Brahmavarchas, editor. Yagyachikitsakebuddhievammedhavardhakprayog. In: YagyaChikitsa, ShriVedmataGayatri Trust, Shantikunj, Haridwar (Uttarakhand), 249411, India; 2012. P. 223-224.
- 46. Babar RP ,Gund SM , Nandgaonkar J. Scientific Evaluation of ApasmarghnaDhoop (Herbal Fumagation) in the Management of Seizure Disorder in Children. International Ayurvedic Medical Journal, 2107; 5 (2):475-481

